

UNCLASSIFIED

PUBLICLY RELEASABLE
LAB. Classification Group

B. Polak

12/13/95

THIS COPY BEARS AUTHORITY FOR
DECLASSIFICATION ACTION
OR FOR UNCLASSIFIED RELEASE
DO NOT CIRCULATE THIS COPY
OUTSIDE OF "D" DIVISION

CLASSIFICATION CANCELLED

For the Atomic Energy Commission

Charles R. Polak
Declassification Officer

2/14/99

**CIC-14 REPORT COLLECTION
REPRODUCTION
COPY**

LAMS-765

Series A

C.3

VERIFIED UNCLASSIFIED

B. Polak

12/13/95

13 August 1948

This document contains 4 pages.

SEARCH FOR SHORT-LIVED Y, Zr, AND Cb

FISSION PRODUCTS

Work done by:

Leon J. Brown

Seymour Katcuff

Report written by:

Leon J. Brown

Seymour Katcuff

LOS ALAMOS NATL. LAB. LIBS.
3 9338 00419 8692

UNCLASSIFIED

UNCLASSIFIED

- 2 -

Search for Short-Lived Y, Zr, and CbFission Products

Table II of "Nuclei Formed in Fission" issued by the Plutonium Project¹ is noteworthy for the great number of gaps it contains where one would expect fission products to exist. In most cases these missing nuclei have not been identified yet because of their short half-lives and the relatively long time required for their chemical isolation. It has recently been shown² that in certain cases the chemical procedures can be shortened by effecting some separation of the fission products from each other during the bombardment by utilizing their different recoil ranges in air. Collimated fission fragments from a thin Pu foil were allowed to pass through air at 140 mm. pressure. By placing a few very thin zapon films at the appropriate distances from the Pu, selected fission fragments could be collected completely separated from fragments of very different masses and partially separated from fragments of neighboring mass numbers.

This procedure is most useful for the separation of yttrium from the rare earth activities since the chemical separation is slow and difficult, and the range method permits a complete and automatic separation. Several runs were made using this technique plus the following chemical operations: one VF_3 precipitation, one $\text{Sr}(\text{NO}_3)_2$ scavenging precipitation from fuming nitric

LOS ALAMOS NATIONAL LABORATORY



3 9338 00419 8692

UNCLASSIFIED

UNCLASSIFIED


- 3 -

acid, another YF_3 precipitation, and a final yttrium oxalate precipitation. Counting was begun as soon as 10 minutes after a 5 minute irradiation. In no case was there a shorter lived component than the previously reported 16.5-min. $Y^{(94)}$. Therefore any short-lived unidentified Y fission products with an appreciable fission yield must have a half-life shorter than 3 minutes.

In the case of zirconium it was found that the range separation technique offered no advantages over the chemical separation from neutron irradiated solutions of uranium or plutonium. In the first few runs, when the standard chemical procedure³ was abbreviated slightly and the final precipitation of zirconium with cupferron was omitted, a new component with a 14-min. half-life appeared in the decay curves. This period was eliminated when four instead of three barium fluozirconate precipitations were performed and when the precipitation with cupferron was included. In the best two runs counting was begun at 15 minutes after the end of 10 minute bombardments. The characteristic growth of 75-min. Cb from 17-hr. Zr was observed but no short-lived activity. Thus no zirconium isotopes with half-lives between 3 minutes and 17 hours result from fission in appreciable yield.

The experiments with columbium were performed both with and without the fission recoil apparatus. The chemical purifications consisted mainly of repeated Cb_2O_5 precipitations and Bi_2S_3


**UNCLASSIFIED**

- 4 -

scavenging precipitations⁴. Small amounts of short period activities were observed when the purely chemical method was used but the half-lives and yields varied from run to run, thus indicating that these were contaminations. In the two runs in which the record method was used counting started at about 16 minutes after 3 minute irradiations. Very little activity was found. Thus any new Cb fission products with an appreciable yield must have shorter half-lives than 4 minutes.

- ¹ J. Am. Chem. Soc. 68, 2411 (1946); Rev. Mod. Phys. 18, 441 (1946).
- ² S. Katcoff, J.A. Miskel, and C.W. Stanley, Report LA-659 (Feb. 1948).
- ³ D.N. Hume, N.E. Ballou, and L.E. Glendenin, Report CN-2815 (June 30, 1945); D.N. Hume, Plutonium Project Record IX B, 8.8.4 (1946).
- ⁴ L. E. Glendenin, Plutonium Project Record IX B, 8.9.1 (1946).

UNCLASSIFIED


THIS COPY BEARS AUTHORITY FOR
DECLASSIFICATION ACTION
OR FOR UNCLASSIFIED RELEASE
DO NOT CIRCULATE THIS COPY
OUTSIDE OF "D" DIVISION

SECRET

SECRET

REC. FILED *Ed. Div*
DATE *8-27-48*
BY

UNCLASSIFIED

