

LA-UR--83-2355

DE83 017321

CONF-8304114--2

Los Alamos National Laboratory is operated by the University of California for the United States Department of Energy under contract W-7405-ENG-36

TITLE: CALCULATION OF NUCLEAR PROPERTIES FAR FROM THE LINE OF β STABILITY

AUTHOR(S): J. Krumlinde, Dept. of Physics and Mathematical Physics, Lund Univ., Sweden
P. Möller, Dept. of Physics and Mathematical Physics, Lund Univ., Sweden
J. R. Nix, T-9

SUBMITTED TO: Contribution to Proc. Workshop on Nuclear Astrophysics, June 6-10, 1983,
Tegernsee, Federal Republic of Germany

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of the authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

By acceptance of this article, the publisher recognizes that the U.S. Government retains a nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes.

The Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy.

MASTER

mg

DISTRIBUTION STATEMENT

Los Alamos National Laboratory
Los Alamos, New Mexico 87545

Contribution to Proc. Workshop on Nuclear Astrophysics,
June 6-10, 1983, at Ringberg castle

Calculation of nuclear properties far from the line of β
stability

J. Krumlinde and P. Möller, Department of Physics and
Mathematical Physics, Lund University, Box 725, S-22007 LUND 7,
Sweden

J. R. Nix, Theoretical Division, Los Alamos National
Laboratory, Los Alamos, New Mexico 87545, USA

July 21, 1983

For a theoretical study of the decay from the r-process line to the line of β stability many nuclear properties must be known or modelled. From the ground state the nucleus may fission or beta decay. In an excited nucleus there is competition between neutron emission, fission and de-excitation by gamma decay. To compute the various decay rates and corresponding decay paths a decay chain program is required. Fission barrier heights, ground state masses and the corresponding particle separation energies are used as input into this decay chain program. Also required is a knowledge of the beta-strength function. At present we have or are developing sophisticated models, based on the microscopic structure of the nuclei involved, to calculate these quantities.

We have, for instance, calculated nuclear potential-energy surfaces, ground-state masses and shapes by use of a Yukawa-plus-exponential macroscopic model and a folded-Yukawa single-particle potential for 4023 nuclei ranging from $^{16}_0$ to $^{279}_{112}$. These calculations, which extend rather far from stability are discussed in detail in refs. ¹⁻³). Here we show in fig. 1 calculated and experimental single particle corrections and the

than 1 MeV. Particularly noteworthy in fig. 1 is that there seems to be no systematic increase in the deviation between calculated and experimental points as you go far from stability, a desirable feature of a mass formula that is to be used for nuclei out to the r-process line. To give specific numbers for a very neutron-rich nucleus we select ^{99}Rb for which the mass excess -50.60 MeV has recently been measured⁴⁾. The present mass formula²⁾ gives -51.66 MeV for this nucleus, a 1.06 MeV error, whereas the droplet model predicts this mass excess to be -56.10 MeV⁵⁾, a 5.5 MeV error. For further discussions we refer to the publications¹⁻³⁾ mentioned above.

We will now briefly discuss the model for the β -strength function we are developing by applying it to the rubidium region. The β -strength function for β^- decay of a sequence of rubidium nuclei to strontium nuclei have been studied experimentally by Kratz and collaborators⁶⁾. They find a pronounced change in appearance of the measured β -strength function between ^{95}Rb and ^{97}Rb . For ^{95}Rb and lighter Rb isotopes the spectra show, in the low-energy region, a very distinct peak structure, with the peaks separated by regions of very low strength. For ^{97}Rb the measured strength function has a much smoother appearance.

These strength functions have been studied in a theoretical model⁷⁾ that takes deformation into account. In the model calculated Nilsson model wave functions, spherical or deformed, as the case may be, are used as the starting point for determining the wave functions of the mother and daughter nuclei in the β decay. Pairing is treated in the BCS approximation. To account for the retardation of low-energy GT-decay rates, a simple residual interaction specific to GT decay, namely $V_{GT} = \chi(\beta^{1-} \cdot \beta^{1+})$, is added to the Hamiltonian, as is customarily done. This residual interaction is treated in the RPA approximation. The strength of the interaction is adjusted to get agreement between the calculated and experimental energy of the giant Gamow-Teller resonance for ^{208}Pb and ^{144}Sm . Since the model is based on calculated wave functions and single-particle levels, studies of nuclei far from stability, where little experimental information is

available, are more straightforward in this model relative to calculations where "experimental" levels are used. The model can treat deformed nuclei employing wave functions calculated to desired accuracy, within the framework of the model, for the deformed single-particle well.

To give an example of results from such a model and to illustrate the effect of deformation on the calculated β^- -strength function we show calculated strength functions for the nucleus ^{97}Rb under the two assumptions of a spherical shape, fig. 2a, or a well-developed deformation, fig. 2b, for the nuclear ground state. The calculation for spherical shape in fig. 2a shows the peak structure so characteristic for the lighter rubidium isotopes, but which is not in agreement with the experimental results for ^{97}Rb . In contrast, the calculation for the deformed case is in excellent agreement with experiment. A more extensive discussion of these results is found in ref.⁷⁾.

This work was supported by the US Department of Energy, the Swedish Natural Science Research Council and NORDITA.

REFERENCES

1. P. Möller and J.R. Nix, Nucl. Phys. A361 (1981) 117
2. P. Möller and J.R. Nix, At. Data and Nucl. Data Tables 26 (1981) 165
3. R. Bengtsson, P. Möller, J.R. Nix and Jing-ye Zhang, to be submitted to Phys. Scr.
4. A. H. Wapstra, 1981 mass evaluation, private communication (1982)
5. W. D. Myers, Droplet Model of Atomic Nuclei (IFI/Plenum, New York, 1977)
6. K. L. Kratz, H. Ohm, A. Schröder, H. Gabelman, W.

Pfeiffer, G. Jung, L. Alquist, G. I. Crawford, Proc. 4th
Int. Conf. on Nuclei far from Stability, Helsingör, 1981,
(CERN 81-09, Geneva, 1981) p. 317

7. J. Krumlinde and P. Möller, submitted to Nucl. Phys.




Figure 1




Figure 2a

