

JUN 6 1988

CONF-880546--10

Los Alamos National Laboratory is operated by the University of California for the United States Department of Energy under contract W-7405-ENG-36

LA--UR--88-1645

DE88 010911

TITLE: IMPROVED CALCULATION OF THE PROMPT FISSION NEUTRON SPECTRUM FROM THE SPONTANEOUS FISSION OF ^{252}Cf

AUTHOR(S): David G. Madland, T-2

SUBMITTED TO: The International Conference on Nuclear Data for Science and Technology, May 30-June 3, 1988, Mito, Japan

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

By acceptance of this article the publisher recognizes that the U S Government retains a nonexclusive, royalty-free license to publish or reproduce the published form of this contribution or to allow others to do so, for U S Government purposes

The Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U S Department of Energy

Los Alamos Los Alamos National Laboratory
Los Alamos, New Mexico 87545

MP

Improved Calculation of the Prompt Fission Neutron Spectrum
From the Spontaneous Fission of ^{252}Cf

D. G. Madland

Los Alamos National Laboratory, Los Alamos, New Mexico, USA

An improved calculation is presented for the prompt fission neutron spectrum $N(E)$ from the spontaneous fission of ^{252}Cf . In this calculation the fission-spectrum model of Madland and Nix is used, but with several improvements leading to a physically more accurate representation of the spectrum. Specifically, the contributions to $N(E)$ from the *entire* fission-fragment mass and charge distributions will be calculated instead of calculating on the basis of a *seven-point approximation* to the peaks of these distributions as has been done in the past.

Therefore, values of the energy release in fission, fission-fragment kinetic energy, nuclear level density, and compound nucleus cross section for the inverse process will be considered on a point-by-point basis over the fragment yield distributions instead of considering averages of these quantities over the peaks of the distributions. Particular attention will be given to the energy-dependent compound nucleus cross sections and to the nuclear level density model. Other refinements to the calculation of $N(E)$ will also be discussed.

Results will be presented and compared with earlier calculations of the spectrum and with recent experimental measurements of the spectrum.

Summary of Theoretical Model

1. Based on standard nuclear evaporation theory
2. Accounts for effects of:
 - motion of fission fragments
 - distribution of fission-fragment excitation energy
 - energy dependence of cross section for inverse process of compound nucleus formation
 - multiple-chance fission effects at high incident neutron energy

3. Original implementation of model:

- **motion of fission fragments:**

calculate *average* kinetic energy per nucleon of the *average* light fragment and likewise for the *average* heavy fragment

$$E_f^L = (A_H/A_L) (\langle E_f^{\text{tot}} \rangle / A), \text{ and}$$

$$E_f^H = (A_L/A_H) (\langle E_f^{\text{tot}} \rangle / A), \text{ where}$$

A_L , A_H , and $\langle E_f^{\text{tot}} \rangle$ are all *average* quantities and A is mass number of fissioning nucleus.

- **distribution of fission-fragment excitation energy:**

$$P(T) = \begin{cases} 2T/T_m^2 & T \leq T_m \\ 0 & T > T_m \end{cases}$$

$$\begin{aligned} \langle E^* \rangle &= \langle E_f \rangle + E_n + B_n - \langle E_f^{\text{tot}} \rangle \\ &= a T_m^2, \text{ where} \end{aligned}$$

$\langle E^* \rangle$ is the total *average* fission-fragment excitation energy,

$\langle E_f \rangle$ is the total *average* energy release in fission,

E_n, B_n are the kinetic and separation energies of the neutron inducing fission,

$\langle E_f^{\text{tot}} \rangle$ is the total *average* fission-fragment kinetic energy, and

a is the nuclear level-density parameter.

The accuracy of T_m therefore depends upon the accuracy of the difference between two large similar numbers, $\langle E_r \rangle$ and $\langle E_f^{\text{tot}} \rangle$, and upon the accuracy of the nuclear level-density parameter a .

We used a seven-point approximation for $\langle E_r \rangle$, experimental values for $\langle E_f^{\text{tot}} \rangle$, and a Fermi gas level-density parameter $a = A/(\text{const})$.

- **energy dependence of cross section for inverse process of compound nucleus formation:**

$\sigma_c(\epsilon)$ calculated on a 100 point grid, to 40 MeV, using the Becchetti-Greenlees potential

omp studies showed that the *shape* of $\sigma_c(\epsilon)$ vs ϵ is the dominant effect

- **multiple-chance fission effects at high incident energy:**

fission probabilities, P_f , from Back and Britt experiments

neutrons emitted prior to fission described by evaporation spectra, including energy-dependent cross sections for the inverse process, $\phi(E, \sigma_c)$

distribution of excitation energy in fissioning nucleus, $D(E^*, \sigma_c)$, given by *complement* of evaporation spectrum $\phi(E, \sigma_c)$

- **basic spectrum in original implementation of model:**

$$N(E) = \frac{1}{2}(N(E, E_f^L, \sigma_C^L) + N(E, E_f^H, \sigma_C^H))$$




Fig. 1.





Fig. 2.



SPONTANEOUS FISSION OF ^{252}Cf

refined implementation of model

- a) include contributions to $N(E)$ from *entire* fission-fragment mass distribution $Y(A)$ instead of the *seven-point approximation*,
- b) include contributions to $N(E)$ from the fission-fragment charge distribution $P(Z)$ for each value of A ,
- c) include contributions to $N(E)$ from the inverse process of compound nucleus formation for the *entire* fission-fragment mass and charge distributions, instead of for the peaks of these distributions,
- d) include explicit gamma-ray de-excitation of fission fragments,
- e) include effects of center-of-mass anisotropy,
- f) investigate scission neutron question.

We have addressed (a), (b), (c), and (f) in the present work

basic spectrum in refined model

$$N(E) = \sum_A \frac{\bar{\nu}(A)}{\bar{\nu}_{\text{tot}}} Y(A) \sum_Z P(Z) N(E, E_f(A), \sigma_c(Z, A), T_m(Z, A))$$

where $\bar{\nu}_{\text{tot}} = \sum_A \bar{\nu}(A) Y(A)$ is the total average multiplicity,

$Y(A)$ is the fission-fragment mass distribution,

$P(Z)$ is the fission-fragment charge distribution,
and

A (or A_L and A_H) and Z are *no longer* average quantities.

- fission-fragment charge distribution $P(Z)$

$$P(Z) = \left\{ \frac{1}{\sqrt{c\pi}} \exp\left(-\frac{(Z-Z_p)^2}{c}\right) \right\}, \text{ which is}$$

normalized by $P_{\text{tot}} = \sum_Z P(Z)$, and where the sum over Z is for all contributing fragment yields for a given fragment mass number A .

the most probable charge, Z_p , is obtained using a corrected unchanged charge distribution (UCD) assumption, due to Unik et al.,

$$\left(Z_p^L - \frac{1}{2} \right) / A_L = \left(Z_p^H + \frac{1}{2} \right) / A_H = Z_c / A_c, \text{ and}$$

$$c = 2(\sigma^2 + 1/12), \quad \sigma = 0.40 \text{ (Reisdorf et al.)}$$

- $T_m(Z,A)$

for spontaneous fission,

$$T_m = \{(E_r(Z,A) - E_f^{\text{tot}}(Z,A))/a(A_c)\}^{1/2}, \text{ where}$$

$$E_r(Z,A) = M - M_L(Z_L, A_L) - M_H(Z_H, A_H)$$

is calculated using experimental masses where they exist and otherwise the calculated masses of Möller and Nix (1986),

and experimental values of E_f^{tot} , averaged over Z , are taken from the measurements of Schmitt et al. (1966),

and the level-density parameter $a(A_c)$ in the Fermi gas model is still given by $a=A_c/\text{const}$.

- kinetic energy per nucleon of fission fragments, $E_f(A)$

calculated using experimental values of $E_f^{\text{tot}}(Z,A)$ due to Schmitt et al. (1966) and momentum conservation.

- inverse cross section $\sigma_c(Z,A)$

calculated using the Becchetti-Greenlees potential for *each* fission fragment.

- fission-fragment mass distribution $Y(A)$

experimental values used from the measurements of Schmitt et al. (1966)

- $\bar{\nu}(A)$

experimental values used from the measurements of Walsh and Boldeman(1977).





Fig. 4.



results

We represented the fission-fragment mass and charge distributions using 28 fragments:

- 14 approximately equispaced A values ranging from 88 to 164
- 2 values of Z per A value, the nearest integer values to Z_p

We used

- 28 optical-model calculations of $\sigma_c(\epsilon)$
- 28 calculations of E_r
- 14 calculations of $E_f(A)$
- 14 experimental values of $E_f^{\text{tot}}(A)$
- 14 experimental values of $\bar{\nu}(A)$





Fig. 1.







Fig. 7.



- We found that the *refined* model calculation compared to the *original* model calculation is
 - *harder* at the high energy end,
 - *harder* at the low energy end, and
 - *softer* in the intermediate region.
- We found that the agreement with the experimental spectrum of Poenitz and Tamura (1982) is *improved* with the refined model calculation, for an identical value of the level-density parameter, namely, $a = A/9.15$
- Using this same value of the level-density parameter, we found *unsatisfactory* agreement with the evaluated spectrum of Mannhart (1987).
- Performing a least-squares adjustment to the evaluated spectrum of Mannhart (1987), we determined a level-density parameter $a = A/9.40$. We again found *unsatisfactory* agreement with the evaluated spectrum of Mannhart (1987).

We conclude that

- the fission-fragment mass and charge distribution should be represented more completely
 - more than 28 fragments,
- the effects of center-of-mass anisotropy should be included, and
- explicit accounting of gamma-ray de-excitation effects should be included.

- Fig. 1 Ratio of the previous least-squares adjusted Los Alamos spectrum and the experimental spectrum of Poenitz and Tamura (1982) to the least-squares adjusted Maxwellian spectrum, for $^{252}\text{Cf}(\text{sf})$.
- Fig. 2 Ratio of calculated to experimental integral cross sections for the spontaneous fission of ^{252}Cf as a function of the effective neutron threshold energy for the reaction. The calculated values are obtained using the previous least-squares adjusted Los Alamos spectrum together with ENDF/B-V pointwise cross sections. The experimental values are those of Grundl et al. and Kobayaski et al.
- Fig. 3 Similar to Fig. 2 except that the calculated integral cross sections are obtained using the least-squares adjusted Maxwellian spectrum.
- Fig. 4 Experimental values of the total fission-fragment kinetic energy for $^{252}\text{Cf}(\text{sf})$ as a function of the heavy fragment mass number, determined by Schmitt et al. (1966).
- Fig. 5 Experimental values of the heavy fragment yield for $^{252}\text{Cf}(\text{sf})$ as a function of the heavy fragment mass number, determined by Schmitt et al. (1966).

Fig. 6 Comparison of the previous least-squares adjusted Los Alamos spectrum, based on considerations of the *peaks* of the fission-fragment mass and charge distributions, and the present Los Alamos spectrum, based on considerations of the *entire* fission-fragment mass and charge distributions. The level-density parameter in both calculations is given by $a=A/(9.15 \text{ MeV})$.

Fig. 7 Ratio of the present Los Alamos spectrum and the evaluated spectrum of Mannhart (1987) to a Maxwellian spectrum with $T=1.42 \text{ MeV}$, for $^{252}\text{Cf}(sf)$. The nuclear level-density parameter is given by $a=A/(9.15 \text{ MeV})$.

Fig. 8 Ratio of the present least-squares adjusted Los Alamos spectrum and the evaluated spectrum of Mannhart (1987) to a Maxwellian spectrum with $T=1.42 \text{ MeV}$, for $^{252}\text{Cf}(sf)$. The adjusted nuclear level-density parameter is given by $a=A/(9.40 \text{ MeV})$.