

CIC-14 REPORT COLLECTION
REPRODUCTION
COPY

PUBLICLY RELEASABLE

Per GB, FSS-16 Date: 2-7-96

By JDB, CIC-14 Date: 2-12-96

[REDACTED]

CLASSIFICATION CANCELLED BY AUTHORITY OF THE JOINT CHIEFS OF STAFF
BY THE DECLASSIFICATION COMMITTEE
C.3

VERIFIED UNCLASSIFIED

Per JLW 6-14-79

By JDB 2-12-96

This is a cover sheet for a document that must be kept unattended or locked when not in use. While this document is in use, you have obtained a sizeable amount of information upon which you have a responsibility that the information contained herein is not to be disseminated of the type that a person may have access to in a locked file or cabinet. Please ensure that you have your own copy of this information.

THE ENEMY WANTS THE INFORMATION CONTAINED IN THIS DOCUMENT.
DON'T LET HIM GET IT THROUGH YOUR NEGLIGENCE.

CLASSIFICATION CANCELLED BY AUTHORITY OF THE JOINT CHIEFS OF STAFF BY THE DECLASSIFICATION COMMITTEE

LOS ALAMOS NATIONAL LABORATORY



3 9338 00330 0422

[REDACTED]

CLASSIFICATION CONTINUED BY AUTHORITY
OF THE BOARD OF GOVERNORS
BY THE DEPARTMENT OF ENERGY
COMMITTEE

LAMS - 345

[REDACTED]

February 6, 1946

This document contains 10 pages

SPIN-DEPENDENT PART OF THE NEUTRON - DEUTERIUM CROSS SECTION

by
T. Hall
F. de Hoffmann



LOS ALAMOS NATL. LAB. LIBS.
3 9338 00330 0422

This document contains information that is the property of the United States Government and is loaned to your agency for your information only. It is not to be distributed outside your agency without the express written approval of the originating agency. The transmission or disclosure of this information in any manner to an unauthorized person is prohibited by law.

CLASSIFICATION CONTINUED BY AUTHORITY
OF THE BOARD OF GOVERNORS
BY THE DEPARTMENT OF ENERGY
COMMITTEE

ABSTRACT

A matrix method for the spin-dependent part of the neutron deuterium scattering cross section is worked out in detail. The method follows closely the treatment of Condon and Shortley for the evaluation of the dot product of matrices.

CONFIDENTIAL
BY [REDACTED]
CLASSIFICATION COMMITTEE

CONFIDENTIAL
BY [REDACTED]
CLASSIFICATION COMMITTEE

-3-

SPIN-DEPENDENT PART OF THE NEUTRON - DEUTERIUM CROSS SECTION

We wish to present a method for the spin-dependent part of the scattering cross section for slow neutrons from ortho and para deuterium. To introduce the problem let us quote from a forthcoming paper by Schwinger and Hammermesh¹⁾:

"We wish to calculate the differential cross section for a scattering process in which a neutron with momentum p^0 collides with a D_2 molecule with momentum $-p^0$ in the internal state specified by the vibrational, rotational and spin quantum numbers v, J, S , thereby producing a neutron with momentum p which has been scattered through the angle Θ into the solid angle $d\Omega$, leaving the molecule in the state characterized by $-p, v', J', S'$. The differential cross section, as computed by the Born approximation, is

$$\sigma_{J',v',S'; J,v,S}(\Theta) d\Omega = \frac{16}{25} \frac{p}{p^0} \frac{1}{2(2v+1)(2J+1)} \sum_{m^1, m_J^1, m_S^1} \sum_{m_1, m_J, m_S} \left| \left(\frac{4}{5} \frac{Mv}{2\pi\hbar} \right)^2 U \psi_i \right|^2 d\Omega \quad (1)$$

which differs formally for the corresponding H_2 molecule cross section²⁾ only by the replacement of the numerical factor 4.9, the square of the reduced mass of the neutron - H_2 molecule system, by 16/25, the square of the reduced mass of the neutron - D_2 molecule system. These reduced mass factors arise

1). Schwinger and Hammermesh, the scattering of slow neutrons by ortho and deuterium. To be published in the Physical Review.

2). Schwinger and Teller, Phys Rev 52, 1937, 286-295 equation 21

in calculating the number of final neutron states per unit range of the total energy, and in the value of the neutron flux relative to the molecule."

Now we know that the interaction can be written as ³⁾

$$U = -\frac{2\pi\hbar^2}{M} \left[\left\{ 4a_{3/2} + 2a_{1/2} \right\} + 2(a_{3/2} - a_{1/2}) \sigma_n \cdot (\sigma_1 + \sigma_2) \right] \left[\delta(r_n - r_1) + \delta(r_n - r_2) \right] \\ - \frac{2\pi\hbar^2}{M} \left[2(a_{3/2} - a_{1/2}) \sigma_n \cdot (\sigma_1 - \sigma_2) \right] \left[(r_n - r_1) - (r_n - r_2) \right] \quad (2)$$

where M is the mass of the neutron; $a_{3/2}$ and $a_{1/2}$ are the scattering amplitudes for a neutron - deuteron quadruplet interaction and a neutron - deuteron doublet interaction respectively; σ_1 and σ_2 are the spin vectors of the deuterons; σ_n is the spin vector of the neutron; r_1 , r_2 and r_n are the respective position vectors and δ is the Dirac delta function. Units are such that the spin of the neutron is one half. The nature and use of these vectors is exactly as that given in Condon and Shortley "Theory of Atomic Spectra", Cambridge 1935 (hereafter referred to as C and S) and we follow their method in deriving the interaction matrices.

Let us set

$$\left. \begin{aligned} \alpha &= 4a_{3/2} + 2a_{1/2} \\ \beta &= 2(a_{3/2} - a_{1/2}) \end{aligned} \right\} \quad (3)$$

thus

$$U = -\frac{2\pi\hbar^2}{M} \left[\alpha + \beta \sigma_n \cdot (\sigma_1 + \sigma_2) \right] \left[(r_n - r_1) + (r_n - r_2) \right] \\ - \frac{2\pi\hbar^2}{M} \left[\beta \sigma_n \cdot (\sigma_1 - \sigma_2) \right] \left[\delta(r_n - r_1) - \delta(r_n - r_2) \right] \quad (4)$$

3). One way of deriving this formula is given by Schwinger and Hammermesh and leads to their equation (6). The apparent difference is due to their definition of σ , in particular $\sigma_n = (1/2)\sigma_{\text{Schwinger}}$

CLASSIFIED BY [REDACTED]
 AUTHORITY [REDACTED]
 DATE [REDACTED]

We note that U is divided into a symmetrical and anti-symmetrical part and rewrite U as

$$U = U_1 \delta(\omega) + U_2 \delta(-\omega) \quad (5)$$

Further we may rewrite the summation part of (1) using this symbolism as

$$\sum_{m^1, m_J^1, m_S^1} \sum_{m^1, m_J^1, m_S^1} \left| \psi_f \frac{MV}{2\pi \hbar^2} U \psi_i \right|^2 = A_1 B_1 + A_2 B_2 \quad (6)$$

where

$$\left. \begin{aligned}
 A_1 &= \sum_{m^1, m_S^1} \sum_{m, m_S} \left| (m^1 m_S \ U_1 \ m^1 \ m_S^1) \right|^2 \\
 A_2 &= \text{same with } |U_2|
 \end{aligned} \right\} \quad (7)$$

and

$$\left. \begin{aligned}
 B_1 &= \sum_{m_J} \sum_{m, m_S} \left| \int \psi_f \delta(\omega) \psi_i \ d\tau \right|^2 \\
 B_2 &= \text{same with } \delta(-\omega)
 \end{aligned} \right\} \quad (8)$$

where ψ_f and ψ_i are the final and initial spacial wave functions of the whole system and τ is their coordinate space.

In the following we wish to show the method for evaluating A_1 and A_2 .
 Let us recall that

- m = z component of neutron spin
- m_J = z component of the rotation vector of the molecule
- m_S = z component of $\sigma_1 + \sigma_2$

Our problem is essentially to obtain the elements of the matrices $\sigma_n \cdot (\sigma_1 + \sigma_2)$

[REDACTED]

-6-

and $\sigma_n \cdot (\sigma_1 - \sigma_2)$. We shall perform this by evaluating $\sigma_n \cdot \sigma_1$ first. The evaluation of $\sigma_n \cdot \sigma_2$ is virtually the same problem. To get $\sigma_n \cdot \sigma_1$ instead of summing over m and m_s , we change to a complete set of commuting variables (J, J_1, J_2, J_z) and sum over j, j_1, j_2 where

$$J_1 = \sigma_1 + \sigma_2$$

$$J_2 = \sigma_n$$

$$J = \sigma_1 + \sigma_2 + \sigma_n$$

$$J_z = \text{the } z \text{ component of } J$$

$$j = \text{the remaining variables completing the set}$$

and small letters denote the eigenvalues of these variables. We note that our J_2 and σ_1 now satisfy the conditions for vectors P and Q as given by C and S page 70 where a scheme for the evaluation of P·Q is given.

The matrix elements of P·Q are seen to be evaluated in terms of two quantities of the form $(\text{---} \begin{matrix} \vdots \\ Q \\ \vdots \end{matrix} \text{---})$ and $(\text{---} \begin{matrix} \vdots \\ P \\ \vdots \end{matrix} \text{---})$. The evaluation of these quantities in detail is given by equation C and S 10⁵ 2a and 10⁵ 2b. Their substitution into the evaluation of 12⁵ 2 then gives P·Q.

For a given molecular spin S, the values of J, J₁ and J₂ will determine the role of the second deuteron spin so that we may apply the argument of C and S regarding relative phases (cf C and S, last paragraph of page 69) with the following results: for elements diagonal in j, j₁ and j₂ the value of $\sigma_n \cdot \sigma_1$ and $\sigma_n \cdot \sigma_2$ are the same; elements off the diagonal are the negative of each other. Therefore the diagonal elements of $\sigma_n \cdot (\sigma_1 - \sigma_2)$ are zero and the off-diagonal elements of $\sigma_n \cdot (\sigma_1 - \sigma_2)$ are zero.

This means that collisions in which the total molecular spin changes are due only to interaction terms of the form $\sigma_n \cdot (\sigma_1 - \sigma_2)$ i.e. only due to U₂, the

-7-

antisymmetrical part of the interaction. Thus for the case in which total molecular spin changes $A_1 = 0$ and we need to evaluate only A_2 . Similarly in those cases in which the total molecular spin remains unchanged $A_2 = 0$ and we need to evaluate only A_1 .

The possible values of S for deuterium are 0, 1 and 2. Let us now treat the individual cases by the method outlined above:

I. SYMMETRIC

$A_2 = 0$ and

$$A_1 = \sum_{j, j_z} \sum_{j_1^1, j_2^1} \left| (j, j_z | \alpha + \beta \sigma_n \cdot (\sigma_1 + \sigma_2) | j_1^1 j_2^1 \right|^2 \quad (9)$$

By our previous arguments regarding $\sigma_n \cdot (\sigma_1 + \sigma_2)$ the matrix $\alpha + \sigma_n \cdot (\sigma_1 + \sigma_2)$ is diagonal (α being just a constant). Substitution into the formula for P-Q also shows that two elements with the same j and different j_z are equal. The summation over j_z therefore gives

$$A_1 = \sum_j (2j+1) \left\{ \alpha + \beta \sigma_n \cdot (\sigma_1 + \sigma_2) \right\}^2 \quad (10)$$

In all symmetric cases $(\leftarrow \vdots P \vdots \rightarrow) = 1$ and $(\leftarrow \vdots Q \vdots \rightarrow) = 1/2$

Let us now look at all the possible symmetric cases in detail

Case I-1:

S goes from $0 \rightarrow 0$

The only possible eigenvalues are

$$j = 1/2, j_1 = 0, j_2 = 1/2$$

Substitution of these values in P-Q gives $\sigma_n \cdot (\sigma_1 + \sigma_2) = 0$

Hence

$$A_1 = 2\alpha^2$$

-8-

Case I - 2:S goes from 1 \rightarrow 1

There are two possible sets of eigenvalues

$$j = 1/2, \quad j_1 = 1, \quad j_2 = 1/2$$

Here $\sigma_n \circ (\sigma_1 + \sigma_2) = 1$

or $j = 1 1/2, \quad j_1 = 1, \quad j_2 = 1/2$

Here $\sigma_n \circ (\sigma_1 + \sigma_2) = 1/2$

Hence

$$\underline{\Lambda_1 = 6\alpha^2 + 3\beta^2}$$

Case I - 3S goes from 2 \rightarrow 2

There are two possible sets of eigenvalues

$$j = 1 \ 1/2, \quad j_1 = 2, \quad j_2 = 1/2$$

Here $\sigma_n \circ (\sigma_1 + \sigma_2) = -3/2$

or $j = 2 \ 1/2, \quad j_1 = 2, \quad j_2 = 1/2$

Here $\sigma_n \circ (\sigma_1 + \sigma_2) = 1$

Hence

$$\underline{\Lambda_1 = 10\alpha^2 + 15\beta^2}$$

CONFIDENTIAL - SECURITY INFORMATION
 BY THE DIRECTOR, FEDERAL BUREAU OF INVESTIGATION
 ON THE BASIS OF A DETERMINATION BY AUTHORITY
 OF THE ATTORNEY GENERAL
 DATE 11/17/54

-9-

II Anti-Symmetrical

By analogous arguments to the symmetrical case we obtain $A_1 = 0$ and

$$A_2 = \sum_j (2j+1) \left\{ \beta \sigma_n \cdot (\sigma_1 - \sigma_2) \right\}^2$$

In all anti-symmetrical cases $(\leftarrow \dot{P} \rightarrow)$ has the value $1/2$ and $(\leftarrow \dot{Q} \rightarrow)$ takes on different values as listed in each case.

Case II - 1

S goes from $0 \rightarrow 1$

For this case $(\leftarrow \dot{Q} \rightarrow) = \sqrt{2/3}$

There are two possible sets of eigenvalues

$$j = 1/2, \quad j_1 = 0 \rightarrow 1, \quad j_2 = 1/2$$

Here $\sigma_n \cdot (\sigma_1 - \sigma_2) = \sqrt{2}$

or $j = 1 \ 1/2, \quad j_1 = 0 \rightarrow 1, \quad j_2 = 1/2$

Here $\sigma_n \cdot (\sigma_1 - \sigma_2) = 0$

Hence $A_2 = 4\beta^2$

Case II - 2

S goes from $1 \rightarrow 0$

This is analogous to the $0 \rightarrow 1$ case, and yields

$$A_2 = 4\beta^2$$

-10-

Case II - 3

S goes from 1 → 2

For this case $(-\ddot{Q}-) = \frac{1}{2\sqrt{3}}$

There are two possible sets of eigenvalues

$$j = 1 \frac{1}{2}, \quad j_1 = 1 \rightarrow 2, \quad j_2 = 1 \frac{1}{2}$$

Here $\sigma_n \cdot (\sigma_1 - \sigma_2) = \sqrt{15}$ or $j = 2 \frac{1}{2}, \quad j_1 = 1 \rightarrow 2, \quad j_2 = 1/2$ Here $\sigma_n \cdot (\sigma_1 - \sigma_2) = 0$

Hence

$$A_2 = 5\beta^2$$

The above method is interesting largely because it can be extended to other elements with relative ease.

The calculation of the hydrogen case is almost no extra work once the deuterium case is dealt with. If the interaction is written of the form of equation (4) where α and β simply have a different meaning then the values of A_1 and A_2 just evaluated change only because of a changed $(-\ddot{P}-)$ and $(-\ddot{Q}-)$ in the case of hydrogen.

We are contemplating the extension of this method to CH_4 and possibly NH_3 .

DOCUMENT ROOM

REC. FROM *SA*

DATE FEB 14 1946

REC. NO. REC. ✓

