

UNCLASSIFIED


CIC-14 REPORT COLLECTION
REPRODUCTION
COPY

CLASSIFICATION CANCELLED

For the Atomic Energy Commission 4-13-48

Charles A. Keller

Declassification Officer *By Pdh*



THIS COPY BEARS AUTHORITY FOR
DECLASSIFICATION ACTION
OR FOR UNCLASSIFIED RELEASE
DO NOT CIRCULATE THIS COPY
OUTSIDE OF "D" DIVISION.

PUBLICLY RELEASABLE
LANL Classification Group

M. Pankratz 3/13/96

LA-656

Series A



This document contains 18 pages.

26 January 1948

THERMODYNAMIC FUNCTIONS FOR TRITIUM AND TRITIUM HYDRIDE.
THE EQUILIBRIUM OF TRITIUM AND HYDROGEN WITH TRITIUM
HYDRIDE. THE DISSOCIATION OF TRITIUM AND TRITIUM HYDRIDE.

Work done by:

Report written by:

Wesley M. Jones

LOS ALAMOS NATIONAL LABORATORY
3 9338 00349 9331

This document contains restricted data within the
meaning of the Espionage Act, and/or
information the disclosure of which in the interest of the
United States would be injurious to the national defense.
The transmission or revelation of its contents in any
manner to an unauthorized person is prohibited and
may result in severe criminal penalties.



UNCLASSIFIED



021154

UNCLASSIFIED


Acknowledgments

The computations for this report were done by
W. M. Jones of CMR-4 and by the following members of
Group T-2:

Bengt Carlson, Group Leader
Josephine Elliott
Max Goldstein
Margaret Johnson
Nell Lane
Marian Peterson
Alice Snowden



UNCLASSIFIED


UNCLASSIFIEDABSTRACT

The heat capacity, entropy, internal energy, and free energy are calculated to 2500° C. for tritium and tritium hydride. The distribution of tritium between the ortho and para states is considered. The equilibrium constant for the formation of tritium hydride from hydrogen and tritium is calculated from ~~298.1~~⁵⁰ to 2500° K. The dissociation of tritium and tritium hydride into the atoms is discussed. The dissociation equilibria of hydrogen, deuterium, and deuterium hydride have been recalculated. The bearing of the radioactivity of tritium and tritium hydride on the attainment of thermodynamic equilibrium is briefly considered.

UNCLASSIFIED


~~CONFIDENTIAL~~Thermodynamic Functions For Tritium And Tritium Hydride.The Equilibrium of Tritium and Hydrogen With Tritium Hydride. The Dissociation of Tritium and Tritium Hydride.

Accurate calculations of the thermodynamic properties of hydrogen have been carried out by Giauque¹, and similar calculations have been made for deuterium and deuterium hydride by Johnston and Long². The present paper presents corresponding work for tritium and tritium hydride. The values refer to a mole of the ideal gas.

The general methods of such calculations are well known. In the present calculations partition functions and their derivatives have been calculated by direct summation over rotational and vibrational states. The energies of T₂ and HT are representable in the form.

$$E_{J,\nu} = \omega_e(\nu + 1/2) - x_e \omega_e(\nu + 1/2)^2 + [B_e - a_e(\nu + 1/2) + \gamma_e(\nu + 1/2)^2 + \delta_e(\nu + 1/2)^3] J(J + 1) + [D_e + \beta_e(\nu + 1/2)] J^2(J + 1)^2 + F_e J^3(J + 1)^3. \quad (1)$$

The constants are listed in Table I. They were computed from the data for ordinary hydrogen as given by Jeppeson³, who expresses the energy in a form slightly different from Eq. (1), through the theory of isotopic diatomic molecules⁴. The excellent agreement of the constants so calculated

¹ Giauque, J. A. C. S. 52, 4826 (1930)

² Johnston and Long, J. Chem. Phys. 2, 389, 710 (1934)

UNCLASSIFIED

~~CONFIDENTIAL~~

TABLE I
MOLECULAR CONSTANTS OF T₂ AND HT

Constant	T ₂ cm. ⁻¹	HT cm. ⁻¹
ω_e	3608.3	2553.8
$X_e \omega_e$	87.585	43.872
B_e	40.5747	20.3243
α_e	1.67048	0.59222
γ_e	0.030434	0.007636
$-\delta_e$	0.0024	0.00042
$-D_e$	0.02068	0.005190
β_e	0.369×10^{-3}	0.0655×10^{-3}
F_e	1.5378×10^{-5}	0.19328×10^{-5}

UNCLASSIFIED

with the observations of Jeppeson⁵ in the case of HD lend confidence to the present procedure. The agreement in the case of $D_2^{2,6}$ is not quite so good. It is felt that, in general, the calculations are correct to the number of significant figures given except at the highest temperatures where the large values of ν and, particularly, of J needed for the convergence of the summations lie outside the range employed by Jeppeson in fitting the constants of Eq. (1).

The constants in Table I were calculated taking the atomic weights (physical scale) of hydrogen and tritium to be 1.008130 and 3.017050⁷. The physical constants are otherwise those of the International Critical Tables⁸ in order that these calculations may best be utilized in connection with the large body of similar calculations now existing. The experiments of Bloch, et al.⁹ indicate the expected one-half unit of spin for the tritium nucleus.

³ Jeppeson, Phys. Rev. 44, 165 (1933)

⁴ Jevons, Report on Band Spectra of Diatomic Molecules, Cambridge University Press (1932)

⁵ Jeppeson, Phys. Rev. 45, 480 (1934)

⁶ Jeppeson, Phys. Rev. 49, 797 (1937)

⁷ Mattauch, Phys. Rev. 57, 1155 (1940)

⁸ International Critical Tables, Volume I, p. 16

⁹ Bloch, Packard, Graves, and Spence, Phys. Rev. 71, 373(L), 551(L) (1947)

UNCLASSIFIED

UNCLASSIFIED

THE EQUILIBRIUM DISTRIBUTION OF MOLECULAR TRITIUM

AMONG ORTHO AND PARA STATES

Like ordinary hydrogen, molecular tritium exists in ortho and para forms which occupy the odd and even rotational states, respectively, and which have the spin weights 3 and 1, respectively. The fraction of the molecules which is ortho attains the limiting value of $3/4$ at moderate temperatures as may be seen from Table II and from Figure I. The limiting value is attained at lower temperatures than is the case for hydrogen and deuterium.

It is well known that both hydrogen and deuterium show a very slow attainment of ortho-para equilibrium in the absence of catalysts. It seems very likely that the radioactivity of tritium, giving rise to ionization, dissociation, and subsequent recombination, would provide an internal mechanism whereby pure tritium could attain ortho-para equilibrium at a much more rapid rate than pure hydrogen or deuterium. Assuming 10^3 bonds broken per disintegration and random recombination of fragments, the half-time for conversion by this mechanism should be of the order of a few days.

THE HEAT CAPACITY AND ENTROPY

The heat capacity at constant pressure of ortho, para, equilibrium, and normal tritium (room temperature ortho-para distribution) are given in Table III along with the corresponding total entropies which include the nuclear spin contribution of $R \ln 4$. Values are also given for HT. By 175°K. the four

UNCLASSIFIED

UNCLASSIFIED





TABLE II

DISTRIBUTION OF MOLECULAR TRITIUM AMONG ORTHO AND PARA STATES

<u>T, °K</u>	<u>% ortho</u>	<u>T, °K</u>	<u>% ortho</u>
0	0.000	40	66.843
10	2.848	50	71.294
15	16.509	75	74.480
20	33.912	100	74.928
25	47.533	125	74.990
30	56.762	150	74.998
35	62.836	175	75.000



UNCLASSIFIED

UNCLASSIFIED





Fig. 1: Fraction of tritium molecules in odd rotational states as a function of temperature.

UNCLASSIFIED

UNCLASSIFIED



types of T_2 have attained identical heat capacities, and the entropies of normal and equilibrium tritium have become identical. Moreover, above this temperature, the entropies of para and of ortho tritium are less than the entropy of equilibrium or of normal tritium by very nearly $R\ln 4 = 2.754$ e. u. and $R\ln 4/3 = 0.572$ e. u., respectively. This simplifies Table III. Figures 2 and 3 show the internal portion of C_V/R at low temperatures. The characteristic features are owing to rotation.

It is well known that calorimetric measurements performed on H_2 in the absence of a catalyst give rise to an entropy lower than the computed value by $R\ln 4 + 3/4R\ln 3$. These two terms correspond to nuclear spin degeneracy and to failure of ortho-para equilibration to occur in the time of experiment, respectively. This "discrepancy" is well understood and has been considered in detail by Giaouque¹. A similar situation exists for D_2 ^{2,10}.

Granting that the calorimetric measurements could be made, we may speculate on the "discrepancy" which would exist between the experimental and computed entropies for T_2 gas. The term $R\ln 4$ would remain even if measurements could be made at a small fraction of one degree Kelvin. If the estimate of a few days for the half time of the ortho-para conversion is accepted, then the remaining discrepancy might be a fair fraction of the corresponding term in H_2 , $3/4R\ln 3$.

¹⁰ Clusius and Bartholome', Z. physikal. Chem. B30, 237 (1935)

UNCLASSIFIED




TABLE III

HEAT CAPACITIES AND ENTROPIES (INCLUDING NUCLEAR SPIN) OF T₂ AND HT IN CAL. MOLE⁻¹ DEG.⁻¹

T, °K	C _p ^o for tritium				S ^o for tritium				C _p ^o for HT	S ^o for HT
	o-T ₂	p-T ₂	e-T ₂	n-T ₂	o-T ₂	p-T ₂	e-T ₂	n-T ₂		
0	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
10	4.967	4.967	6.770	4.967	18.860	14.495	14.876	18.886	4.977	16.043
15	4.967	4.981	8.970	4.971	20.874	16.510	18.121	20.901	5.143	18.083
20	4.967	5.104	8.689	5.001	22.304	17.956	20.703	22.334	5.611	19.620
25	4.967	5.451	7.745	5.088	23.414	19.127	22.539	23.460	6.199	20.936
30	4.998	5.999	7.043	5.248	24.321	20.167	23.884	24.400	6.680	22.111
35	5.056	6.615	6.633	5.446	25.095	21.138	24.936	25.223	6.974	23.165
40	5.154	7.163	6.423	5.656	25.776	22.059	25.806	25.964	7.111	24.107
50	5.461	7.992	6.393	6.094	26.957	23.737	27.224	27.270	7.141	25.701
75	6.335	7.657	6.657	6.665	29.344	26.831	29.848	29.833	7.023	28.571
100	6.787	7.144	6.876	6.876	31.240	29.004	31.798	31.798	6.987	30.585
125	6.925	7.007	6.945	6.945	32.772	30.580	33.341	33.341	6.977	32.143
150	6.958	6.975	6.963	6.963	34.038	31.854	34.610	34.610	6.974	33.415
175	6.966	6.970	6.967	6.967	35.112	32.929	35.683	35.683	6.973	34.490

T, °K	C _p ^o	S ^o	C _p ^o	S ^o	T, °K	C _p ^o	S ^o	C _p ^o	S ^o
	T ₂	e-T ₂	HT	HT		T ₂	e-T ₂	HT	HT
200	6.969	36.614	6.973	35.421	700	7.347	45.443	7.097	44.186
225	6.971	37.435	6.974	36.242	800	7.512	46.434	7.182	45.139
250	6.973	38.169	6.975	36.977	900	7.676	47.329	7.286	45.991
275	6.975	38.834	6.976	37.642	1000	7.832	48.146	7.403	46.764
298.1	6.978	39.397	6.978	38.205	1250	8.168	49.931	7.710	48.449
300	6.978	39.441	6.978	38.249	1500	8.427	51.445	7.995	49.881
400	7.007	41.452	6.986	40.257	2000	8.781	53.921	8.444	52.247
500	7.078	43.022	7.002	41.818	2500	9.013	55.907	8.760	54.166
600	7.197	44.322	7.037	43.097					

APPROVED FOR PUBLIC RELEASE

APPROVED FOR PUBLIC RELEASE

UNCLASSIFIED

UNCLASSIFIED




Fig. 2: Internal portion of the low temperature

UNCLASSIFIED




Fig. 3: Internal portion of the low temperature heat capacities at constant volume of equilibrium tritium, normal tritium, and tritium hydride.

UNCLASSIFIED

THE INTERNAL ENERGY AND THE FREE ENERGY

The internal energy and the free energy, both given with reference to the state in which $\nu = J = 0$ are given in Tables IV and V. Table IV does not include the contribution $(E - E_0^{\circ})_{\text{translation}} = 3/2 RT$. The entries of Table V include the nuclear spin contribution of $R \ln 4$ for T_2 and HT and of $R \ln 4$ for atomic T, arising from the half units of nuclear and electron spin.

THE EQUILIBRIUM OF HYDROGEN AND TRITIUM WITH TRITIUM HYDRIDE

Values of $\left(\frac{F^{\circ} - E_0^{\circ}}{T}\right)$ for H_2 , T_2 and HT may be employed to calculate the equilibrium constant, K, for the reaction $H_2 + T_2 = 2HT$ from the relationship:

$$-R \ln K = \Delta \left(\frac{F^{\circ} - E_0^{\circ}}{T} \right) - \frac{\Delta E_0^{\circ}}{T} \quad (2)$$

The values of $\left(\frac{F^{\circ} - E_0^{\circ}}{T}\right)$ for H_2 were taken from the tables of Giauque¹ at 298.1° K. and at higher temperature, while the values of Urey and Rittenberg¹¹, corrected to $R = 1.9869 \text{ cal. deg.}^{-1} \text{ mole}^{-1}$, were used below this temperature. The quantity ΔE_0° is obtained from the zero point energies 6184.3, 3597.9, and 5065.0 cal. mole⁻¹ for H_2 , T_2 and HT, and it has the value 347.8 cal. mole⁻¹. The possibility of isotopic electronic shifts^{2,5,6} has not been considered in this and the following section. Table VI gives values of K at several temperatures.

UNCLASSIFIED

UNCLASSIFIED

TABLE IV

E - E₀⁰ IN CAL. DEG.⁻¹ MOLE⁻¹ FOR T₂ AND HT

(EXCLUDING TRANSLATION)

T, °K	o-T ₂	p-T ₂	e-T ₂	n-T ₂	HT
0	113.79	0.00	0.00	85.34	0.00
10	113.79	0.00	3.24	85.34	0.01
15	113.79	0.02	18.80	85.35	0.35
20	113.79	0.32	38.80	85.42	2.29
25	113.84	1.77	55.02	85.82	6.98
30	113.89	5.50	67.02	86.80	14.42
35	114.17	12.20	76.27	88.68	23.79
40	114.84	21.86	84.01	91.60	34.22
50	118.14	47.51	97.87	100.49	55.97
75	141.90	113.02	135.74	134.69	108.67
100	182.68	176.66	181.17	181.18	159.52
125	230.27	228.98	229.95	229.95	209.87
150	279.73	279.48	279.67	279.67	260.07
175	329.63	329.58	329.62	329.62	310.22
200	379.65	379.64	379.65	379.65	360.36

T, °K	T ₂	HT	T, °K	T ₂	HT
225	429.72	410.52	700	1436.6	1379.9
250	479.84	460.70	800	1682.8	1596.9
275	530.01	510.91	900	1945.5	1823.5
298.1	576.43	557.33	1000	2224.3	2061.1
300	580.25	561.15	1250	2984	2708
400	782.49	762.56	1500	3818	3431
500	989.6	965.1	2000	5644	5063
600	1206.3	1170.1	2500	7612	6885

UNCLASSIFIED



TABLE V

$-\left(\frac{F^{\circ} - E_0^{\circ}}{T}\right)$ FOR $e-T_2$, HT, and T IN CAL. DEG.⁻¹ MOLE⁻¹
(SPIN EFFECTS INCLUDED)

T, °K	$e-T_2$	HT	T, °K	$e-T_2$	HT	T
10	9.585	11.075	250	31.283	30.167	26.205
15	11.900	13.092	275	31.940	30.817	26.679
20	13.795	14.539	298.1	32.496	31.368	27.080
25	15.371	15.689	300	32.540	31.411	27.111
30	16.683	16.663	400	34.528	33.384	28.540
35	17.789	17.518	500	36.076	34.921	29.649
40	18.739	18.284	600	37.344	36.179	30.554
50	20.299	19.614	700	38.423	37.248	31.320
75	23.071	22.155	800	39.364	38.175	31.983
100	25.019	24.023	900	40.200	38.997	32.563
125	26.535	25.497	1000	40.955	39.736	33.092
150	27.778	26.714	1250	42.576	41.315	34.195
175	28.833	27.750	1500	43.932	42.627	35.106
200	29.748	28.652	2000	46.132	44.748	36.535
225	30.558	29.451	2500	47.895	46.445	37.643

UNCLASSIFIED







TABLE VI
EQUILIBRIUM CONSTANT FOR $H_2 + T_2 = 2HT$

<u>T, °K</u>	<u>K, [REDACTED]</u>	<u>T, °K</u>	<u>K, [REDACTED]</u>
50	0.243	700	3.59
100	0.947	800	3.69
200	1.953	900	3.76
298.1	2.58	1000	3.82
300	2.59	1250	3.91
400	2.99	1500	3.95
500	3.26	2000	3.99
600	3.45	2500	4.00

UNCLASSIFIED



THE DISSOCIATION OF ISOTOPIC HYDROGEN MOLECULES **UNCLASSIFIED**

The dissociation constants of T_2 and HT have been calculated from a relationship of the form of Eq. (2). The value of ΔE_0 is found from $D_0(H_2)$ and the difference in the zero point energies of H_2 and the isotopic molecule. The work of Beutler and Jünger¹² provides the accurate value of $103,222 \pm 17$ cal. mole⁻¹ for $D_0(H_2)$. The values for the dissociation constants of H_2 , HD and D_2 ^{1,2} have been corrected using the present value of $D_0(H_2)$. Dissociation constants are presented in Table VII. The corresponding percentages of dissociation for a total pressure of one atmosphere are given in Table VIII.

¹¹ Urey and Rittenberg, J. Chem. Phys. 1, 137 (1933)

¹² Beutler and Jünger, Z. Physik 101, 304 (1936)

UNCLASSIFIED

UNCLASSIFIED

TABLE VII
DISSOCIATION CONSTANTS OF ISOTOPIC HYDROGEN MOLECULES
IN ATMOSPHERES

T, °K	1000	1500	2000	2500
H ₂	5.22 x 10 ⁻¹⁸	3.13 x 10 ⁻¹⁰	2.66 x 10 ⁻⁶	6.35 x 10 ⁻⁴
HD	2.00	1.36	1.23	3.01
D ₂	3.00	2.34	2.24	5.67
HT	1.82	1.30	1.19	2.94
T ₂	2.44	2.12	2.11	5.44

TABLE VIII


PERCENTAGE DISSOCIATION OF ISOTOPIC HYDROGEN MOLECULES
AT A TOTAL PRESSURE OF ONE ATMOSPHERE

T, °K	1000	1500	2000	2500
H ₂	11 x 10 ⁻⁸	8.8 x 10 ⁻⁴	8.2 x 10 ⁻²	1.26
HD	7.1	5.8	5.5	0.87
D ₂	8.7	7.7	7.5	1.19
HT	6.8	5.7	5.4	0.86
T ₂	7.6	7.3	7.3	1.17

UNCLASSIFIED

UNCLASSIFIED

UNCLASSIFIED



THIS COPY BEARS AUTHORITY FOR
DECLASSIFICATION ACTION
OR FOR UNCLASSIFIED RELEASE
DO NOT CIRCULATE THIS COPY
OUTSIDE OF "D" DIVISION

DOCUMENT ROOM

REC. FROM Ed. Div
DATE 2-4-48
REC. NO. REC.

96109

UNCLASSIFIED

UNCLASSIFIED