


LA-4851

CIC-14 REPORT COLLECTION
**REPRODUCTION
COPY**

C. 3

$\text{Be}^9(d,n)\text{B}^{10}$ Cross Section in the Range
0 to 2.8 MeV



los alamos
scientific laboratory

of the University of California

LOS ALAMOS, NEW MEXICO 87544



This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

Printed in the United States of America. Available from
National Technical Information Service
U. S. Department of Commerce
5285 Port Royal Road
Springfield, Virginia 22151
Price: Printed Copy \$3.00; Microfiche \$0.95

LA-4851

UC-34

ISSUED: May 1972



los alamos
scientific laboratory
of the University of California
LOS ALAMOS, NEW MEXICO 87544

$\text{Be}^9(d,n)\text{B}^{10}$ Cross Section in the Range 0 to 2.8 MeV



by

J. J. Devaney



$\text{Be}^9(d,n)\text{B}^{10}$ CROSS SECTION IN THE RANGE
0 TO 2.8 MeV

by

J. J. Devaney

ABSTRACT

The total $\text{Be}^9(d,n)\text{B}^{10}$ cross section is constructed by theoretically completing an experimental partial absolute cross section and using it to normalize a relative measurement which in turn was extrapolated to below 0.5 MeV by the relative energy dependence of the coulomb penetration factor.

As a result of recent interest in the $\text{Be}^9(d,n)$ reaction as a neutron source, we have determined its cross section for total neutron production in the 0- to 2.8-MeV range. We were able to find partial absolute cross sections¹ at 1.1 MeV and above, to which we add theoretical estimates of the remaining partial cross sections achieving thereby the total (d,n) cross section. This result was used to normalize a relative measurement² that, in turn, was extended to low energies by the relative energy dependence of the coulomb penetration factor.

RESULTS

The $\text{Be}^9(d,n)\text{B}^{10}$ cross section is given by Eq. (1) for energies $E \leq 0.2$ MeV, and in Table I for $E = 0.05$ to 2.785 MeV, all in millibarns.

$$\sigma \approx \frac{1.94 \times 10^5}{E} \cdot e^{-5.965E^{-1/2}} \quad (\text{mb}) \quad (1)$$

E in MeV and $E \leq 0.2$ MeV.

The relative error of the Koltay versus the Siemssen et al. results at 1.1 and 1.55 MeV is $\pm 12\%$. To that one must add the experimental error of the relative point² at 0.5 MeV that is used to normalize the lower energy coulomb extrapolation, namely 26%. In addition, consistency checks among experiments, 0.5 to 0.6 MeV, and several coulomb extrapolations lead to confidence levels of about $\pm 5\%$ in the transition from 0.5 MeV to lower energies. On the other end of the energy range, our highest point (2.785

MeV) may be taken to have a theoretical probable error of about 16 mb out of 428 mb, (say we missed a spin assignment or a transmission by a factor of 2), which is 3.7%, with experimental error in the 20% neighborhood. Consequently, these results must certainly be regarded as approximate and errors of the order of 29% cannot be excluded.

TABLE I
 $\text{Be}^9(d,n)\text{B}^{10}$ CROSS SECTION

E MeV	σ (mb)
0.05	1×10^{-5}
0.1	0.0125
0.2	1.56
0.3	12.5
0.4	38.7
0.5	77.7
0.6	112
0.7	158
0.8	217
0.9	298
1.0	348
1.05	364
1.1	337
1.2	280
1.3	236
1.37	228
1.4	230
1.5	283
1.6	384
1.95	413
2.37	433
2.785	428

CALCULATIONAL METHOD

From Siemseen, Cosack, and Felst¹ we have an absolute measurement of $\sigma[\text{Be}^9(d,n)\text{B}^{10}, \text{B}^{10*}]$ to the first five levels of B^{10} , including ground state, from 1.1 to 2.785 MeV laboratory deuteron energy. These levels we label 0, 1, 2, 3, 4 and they are given in Table III. Table II (taken from Ref. 3) gives the character of the levels.

TABLE II
LEVELS OF B^{10}

Level Number	Energy (MeV)	J^π, T	$Q \text{Be}^9(d,n)$ (MeV)(Ref.4)	Reaction Statistical Wt for l
0	0	3+,0	4.363	
1	0.7173	1+,0	3.6457	
2	1.74	0+,1	2.623	
3	2.154	1+,0	2.209	
4	3.585	2+,0	0.778	$l=1:0.125$
5	4.774	(2+),0	-0.411	as level 4
6	5.114	(2-),0	-0.751	$l=0:.0167;2:.25$
7	5.164	2+,1	-0.801	as level 4
8	5.183	1+,0	-0.820	$l=1:0.208$
9	5.923	2+,0	-1.560	as level 4
10	6.029	4+	-1.666	$l=1:0.0139$
11	6.133	2+assumed	-1.770	as level 4
12	6.566	2+assumed	-2.203	as level 4

TABLE III
 $\sigma[\text{Be}^9(d,n)\text{B}^{10}, \text{B}^{10*}]^1$
IN MILLIBARNS

To Level of B^{10}	0	1	2	3	4
Deuteron Energy (MeV)					
1.1	81	95	13	49	57
1.55	82	107	12	58	72
1.95	91.5	108.5	12	54	61
2.37	85.5	99.7	11.8	49	64.1
2.785	76	83	11.8	40.7	60.5

TABLE IV

To Level B^{10}	$\sigma[\text{Be}^9(d,n)\text{B}^{10*}]$ IN MILLIBARNS									
Deuteron Energy (MeV)	5	6	7	8	9	10	11	12	Sum 0 to 12	
		(Sum $l=0,2$)								
1.1	14.2	7.2	1.3	1.3	0	0	0	0		319
1.55	30.1	13.4	13.5	13.5	0	0	0	0		401.5
1.95	32.5	13.1	20.	19.9	0.3	0	0	0		413
2.37	40.0	15.6	28.8	28.7	7.2	.5	2.5	0		433
2.785	41.9	16.6	32.8	32.8	14.3	1.3	14.8	1.		428

For a precise calculation of the contributions to levels 5 through 12 we need the relevant compound nucleus theory plus stripping theory with the associated input parameters.⁵ In the absence of these and in keeping with the accuracy and paucity of the

data we have, it is appropriate to use simpler methods. Accordingly we calculate the cross section to the higher levels by multiplying the known cross section to level four by penetrability and statistical weight ratios between level four and the desired level. Lacking intermediate state spin and orbital parameters and because of the low deuteron energy, we take the incoming deuteron orbital angular momentum to be zero, and then weight every possible resulting angular momentum path equally, which procedure is really accurate on the average only. In our energy range for the final states of level 4 and above, we need only consider outgoing $l = 0, 1, 2$ because the penetration of higher l channels is quite negligible to our accuracy. The reaction statistical weight found therefore by simple counting is listed in Table II, fifth column. For levels 5, 7, 9, 11, and 12 the statistical factor drops out because they have the same spin, parity, and isospin as level 4, (2+,0), so statistical errors to these levels may be considered of a lower order, or zero.

For the relative angular momentum barrier penetration we use the simple formulas of Blatt and Weisskopf⁶ with a well depth of (49.3-0.33E)MeV and an effective radius of 2.7 fermis.⁷ Table IV shows the partial cross section to the higher excited states of B^{10} thus obtained. Table IV also gives the total (d,n) cross section from levels 0 to 12 of Tables III and IV.

We use the results at 1.1 and 1.55 MeV weighted 0.8 and 0.2 respectively, to normalize Koltay's relative curve² which with Table IV gives the cross-

section from 0.5 to 2.785 MeV. For the remainder we assume that the dominant energy dependence is that of the coulomb barrier penetration, which has been verified for the simple formula (1) up to 0.25 MeV (Ref.8) and for a WKBJ coulomb penetration up to

0.9 (Ref. 2). We use a more accurate machine calculation of the coulomb transmission for a $\text{Be}^9\text{-d}$ reaction radius of 7 fermis, provided for us by G. M. Hale⁹, to extrapolate the cross section from 0.5 to 0.2 MeV and then use Eq. (1) for the lower energies.

ACKNOWLEDGMENT

We take pleasure in acknowledging the kindness of Drs. F. K. McGowan of Oak Ridge, J. A. Phillips and D. G. Foster, Jr. of LASL, who furnished relevant portions of their work, to D. R. Harris, Jr. and J. Carson Mark for critical reviews, to D. C. Dodder for a helpful discussion, and to G. M. Hale for coulomb penetration ratios.

REFERENCES

1. R. H. Siemssen, M. Cosack, and R. Felst, "Reaction Mechanism Study of $L_p = 1$ Stripping Processes on Nuclei of the $1p$ Shell," Nucl. Phys. 69, 209-226 (1965).
2. E. Koltay, "Excitation Function of the Nuclear Reaction ${}^9\text{Be}(d,n){}^{10}\text{B}$ by Artificially Accelerated Particles in the 0.5-1.6 MeV. Energy Range," Acta Phys. Acad. Sci. Hung. 16, 93 (1963).
3. T. Lauritsen and F. Ajzenberg-Selove, "Energy Levels of Light Nuclei, VII. A=5-10," Nucl. Phys. 78, 1-176 (1966).
4. F. Ajzenberg-Selove and T. Lauritsen, "Energy Levels of Light Nuclei, VII. A=11-12," Nucl. Phys. A114, 1-142 (1968).
5. A. I. Shpetnyi, "Energy and Angular Distribution of Neutron Emitted in the $\text{Be}^9(d,n)\text{B}^{10}$ Reactions," Soviet Phys. JETP 5, 357-364 (1957).
6. J. M. Blatt and V. F. Weisskopf, Theoretical Nuclear Physics (Wiley, New York, 1952).
7. F. P. Agee and L. Rosen, "Calculations of Neutron Cross Sections Using a Local Optical Potential with Average Parameters," Los Alamos Scientific Laboratory report LA-3538-MS, Vol. 1, (September 1966).
8. J. A. Phillips and G. A. Sawyer, "Survey of Several of the Reactions Occurring when Lithium and Beryllium are Bombarded by Protons and Deuterons of 30-250 KeV," Los Alamos Scientific Laboratory report LA-1578 (August 1953). Verified for $\text{Be}^9(d,n)$ by D. C. Ralph and F. E. Dunnam, "Neutron Yields as a Function of Energy in the Reactions $\text{Li}^7(d,n)\text{Be}^8$ and $\text{Be}^9(d,n)\text{B}^{10}$," Phys. Rev. 98, 249 (1955).
9. G. M. Hale, Los Alamos Scientific Laboratory, private communication, March 1972.